arXiv:1302.2354v2 [math.MG] 12 Feb 2013

A SHORT PROOF OF KLEE'S THEOREM

JOHN ZANAZZI, NORTHERN ARIZONA UNIVERSITY

ABSTRACT. In 1959, Klee proved that a convex body K is a polyhedron if and only if all of its projections are polygons. In this paper, a new proof of this theorem is given.

1. INTRODUCTION

In this note we give a shorter proof of the following theorem in [1].

Theorem 1.1. *Let K be a convex body in $\mathbb{R}^3$ such that any orthogonal projection to a plane is a polygon. Then K is a polyhedron.*

Klee's proof is based on [2] which states that a convex closed cone is polyhedral if and only if all its projections are closed. Our proof uses properties of the polar dual.

2. DUAL REFORMULATION

Recall that a *convex body* is a closed bounded convex set with nonempty interior.

Fix a convex body K in $\mathbb{R}^3$. Assume that the origin of $\mathbb{R}^3$ belongs to the interior of K.

The *polar dual* body of K will be denoted as K^*; i.e.,

$$K^* = \{\, y \mid x \cdot y \leq 1 \text{ for any } x \in K \,\}$$

Clearly K^* is a convex body and the origin is an interior point of K^*. Moreover K^* is a convex polyhedron if and only if so is K.

Let P be a plane in $\mathbb{R}^3$. The orthogonal projection to P will be denoted as π_P.

The following statement follows directly from the definition of polar dual.

Proposition 2.1. *Let P be a plane passing through the origin. Then*

$$K^* \cap P = \pi_P(K)^* \cap P.$$

Note that $\pi_P(K)^* \cap P$ is polygon if and only if so is $\pi_P(K)$. Using the above proposition the Klee theorem can be reformulated the following way.

Theorem 2.2. *Let K^* be a convex body in $\mathbb{R}^3$ which contains the origin in its interior. Assume that for any plane P passing through the origin the intersection $P \cap K^*$ is a polygon. Then K^* is a polyhedron.*

3. PROOF OF THEOREM 2.2

Let us denote by ∂K^* the boundary of K^*.

Lemma 3.1. *Let K^* be a convex body in $\mathbb{R}^3$ and $p, q, x, y \in K^*$ be points such that x lies between p and q. Assume that the line segment $[xy]$ lies completely in ∂K^*. Then the triangle $\triangle pqy$ lies completely in ∂K^*.*



FIGURE 1.

Proof. Note that the midpoint of $[xy]$ lies in the interior of $\triangle pqy$.

Further note that if one point in $\triangle pqy$ belongs to the interior of K^* then so are all the interior points of $\triangle pqy$. Hence the result follows. □

Proposition 3.2. *Assume K^* be as in Theorem 2.2. Let $p,q \in K$. Then there exists $\varepsilon > 0$ such that for $x \in K$, if $0 < |p-x| < \varepsilon$ and $\angle xpq < \varepsilon$, then x is not extreme.*

Proof. The statement is evident if $[pq]$ passes through the interior of K^*, so we can assume that $[pq] \subset \partial K^*$.

Denote by x the midpoint of $[pq]$. Choose a plane P through the origin which intersects $[pq]$ transversely at x.

The intersection $K^* \cap P$ is a polygon, where the sides extending from x are denoted by $[xy]$ and $[xz]$. Refer to Figure 1 for clarity.

Applying Lemma 3.1 we get that the triangles $\triangle pyq$ and $\triangle pzq$ lie completely in ∂K^*. Hence the result follows. □

Proof of Theorem 2.2. Assume K^* has infinite sequence of distinct extreme points $q_1, q_2, \ldots$.

We may pass to a convergent subsequence q_n which has the property that the unit vectors $v_n = \frac{q_n - p}{|q_n - p|}$ also converge, say $v_n \to u$. Let $p \in \partial K$ be the point such that $q_n \to p$ as $n \to \infty$.

Consider the plane P which passes through p, u and the origin. Since the intersection $P \cap K^*$ is a polygon, there is a line segment $[pq] \subset \partial K^*$ which goes from p in the direction of u.

Applying Proposition 3.2, we arrive to a contradiction. □

4. ACKNOWLEDGMENTS

I am greatly indebted to Anton Petrunin, Greg Kuperberg, and Sergei Tabachnikov for suggestions and editing drafts. This paper would not have been possible without them.

REFERENCES

[1] V. Klee. Some characterizations of convex polyhedra. *Acta Mathematica*, pages 79-107, 1959.
[2] H. Mirkil. New characterizations of polyhedral cones. *Canadian Journal of Mathematics*, pages 1-4, 1957.